

# Numerical Analysis - Part II

Anders C. Hansen

Lecture 19

---

*Iterative methods for linear algebraic systems*

# Minimization of quadratic function

The methods we considered so far for solving  $A\mathbf{x} = \mathbf{b}$ , namely Jacobi, Gauss-Seidel, and those with relaxation, fit into the scheme

$$\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + c_k \mathbf{d}^{(k)},$$

where we were aimed at getting  $\rho(H) < 1$  for the iteration matrix  $H$ . Say, for Jacobi with relaxation, we set  $c_k = \omega$  and  $\mathbf{d}^{(k)} = D^{-1}(\mathbf{b} - A\mathbf{x}^{(k)})$ .

For solving  $A\mathbf{x} = \mathbf{b}$  with a (positive definite) matrix  $A > 0$ , there is a different approach to constructing good iterative methods. It is based on successive minimization of the quadratic function

$$F(\mathbf{x}^{(k)}) := \|\mathbf{x}^* - \mathbf{x}^{(k)}\|_A^2 = \|\mathbf{e}^{(k)}\|_A^2,$$

since the minimizer is clearly the exact solution. Here,  $\|\mathbf{y}\|_A := (A\mathbf{y}, \mathbf{y})^{1/2} := \sqrt{\mathbf{y}^T A \mathbf{y}}$  is a Euclidean-type distance which is well-defined for  $A > 0$ .

# Minimization of quadratic function

---

So, at each step  $k$ , we are decreasing the  $A$ -distance between  $\mathbf{x}^{(k)}$  and the exact solution  $\mathbf{x}^*$ . Thus, for a symmetric positive definite  $A > 0$ , we choose an iterative method that provides the descent condition

$$\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + c_k \mathbf{d}^{(k)} \quad \Rightarrow \quad F(\mathbf{x}^{(k+1)}) < F(\mathbf{x}^{(k)}). \quad (1)$$

# Minimization of quadratic function

---

An equivalent approach is to minimize the quadratic function

$$F_1(\mathbf{x}) = \frac{1}{2}\mathbf{x}^T A\mathbf{x} - \mathbf{x}^T \mathbf{b},$$

which attains its minimum when  $\nabla F_1(\mathbf{x}) = A\mathbf{x} - \mathbf{b} = 0$ , and which does not involve the unknown  $\mathbf{x}^*$ . It is easy to check that  $F_1(\mathbf{x}) = \frac{1}{2}F(\mathbf{x}) - \frac{1}{2}c$ , where  $c = \mathbf{x}^{*T}A\mathbf{x}^*$  is a constant independent of  $k$ , hence equivalence.

## Quadratic function – Jacobi and Gauss–Seidel

---

Both the Jacobi and the Gauss–Seidel methods satisfy (1), precisely

$$(A\mathbf{e}^{(k+1)}, \mathbf{e}^{(k+1)}) = (A\mathbf{e}^{(k)}, \mathbf{e}^{(k)}) - (C\mathbf{y}^{(k)}, \mathbf{y}^{(k)}) < (A\mathbf{e}^{(k)}, \mathbf{e}^{(k)}),$$

where for Gauss-Seidel:  $C = D > 0$ ,  $\mathbf{y}^{(k)} := (L_0 + D)^{-1}A\mathbf{e}^{(k)}$ ;

and for Jacobi:  $C = 2D - A > 0$ ,  $\mathbf{y}^{(k)} := D^{-1}A\mathbf{e}^{(k)}$ .

# A-orthogonal projection

---

**A-orthogonal projection method:** Next, we strengthen the descent condition (1), namely given  $\mathbf{x}^{(k)}$  and some  $\mathbf{d}^{(k)}$  (called a *search direction*), we will seek  $\mathbf{x}^{(k+1)}$  from the set of vectors on the line  $\ell = \{\mathbf{x}^{(k)} + \alpha \mathbf{d}^{(k)}\}_{\alpha \in \mathbb{R}}$  such that it makes the value of  $F(\mathbf{x}^{(k+1)})$  not just smaller than  $F(\mathbf{x}^{(k)})$ , but as small as possible (with respect to this set), namely

$$\mathbf{x}^{(k+1)} := \arg \min_{\alpha} F(\mathbf{x}^{(k)} + \alpha \mathbf{d}^{(k)}). \quad (2)$$

## Lemma 1

*The minimizer in (2) is given by the formula*

$$\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha_k \mathbf{d}^{(k)}, \quad \alpha_k = \frac{(\mathbf{r}^{(k)}, \mathbf{d}^{(k)})}{(A\mathbf{d}^{(k)}, \mathbf{d}^{(k)})}. \quad (3)$$

This choice of  $\alpha_k$  is referred to as exact line search.



# A-orthogonal projection

**Proof.** From the definition of  $F$ , it follows that in (2) we should choose the point  $\mathbf{x}^{(k+1)} \in \ell$  that minimizes the  $A$ -distance between  $\mathbf{x}^*$  and the points  $\mathbf{y} \in \ell$ . Geometrically, it is clear that the minimum occurs when  $\mathbf{x}^{(k+1)}$  is the  $A$ -orthogonal projection of  $\mathbf{x}^*$  onto the line  $\ell = \{\mathbf{x}^{(k)} + \alpha \mathbf{d}^{(k)}\}$ , i.e., when

$$\begin{aligned}\mathbf{x}^* - \mathbf{x}^{(k+1)} \perp_A \mathbf{d}^{(k)} &\Rightarrow A(\mathbf{x}^* - \mathbf{x}^{(k+1)}) \perp \mathbf{d}^{(k)} \\ &\Rightarrow \mathbf{r}^{(k+1)} = \mathbf{r}^{(k)} - \alpha_k A \mathbf{d}^{(k)} \perp \mathbf{d}^{(k)}.\end{aligned}$$

This gives expression for  $\alpha_k$  in (3). □

# The steepest descent method


**The steepest descent method:** This method takes  $\mathbf{d}^{(k)} = -\nabla F_1(\mathbf{x}^{(k)}) = \mathbf{b} - A\mathbf{x}^{(k)}$  for every  $k$ , the reason being that, locally, the negative gradient of a quadratic function shows the direction of the (locally) steepest descent at a given point. Thus, the iterations have the form

$$\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha_k(\mathbf{b} - A\mathbf{x}^{(k)}), \quad k \geq 0. \quad (4)$$


It can be proved that the sequence  $(\mathbf{x}^{(k)})$  converges to the solution  $\mathbf{x}^*$  of the system  $A\mathbf{x} = \mathbf{b}$  as required, but usually the speed of convergence is rather slow.

The reason is that the iteration (4) decreases the value of  $F(\mathbf{x}^{(k+1)})$  locally, relatively to  $F(\mathbf{x}^{(k)})$ , but the global decrease, with respect to  $F(\mathbf{x}^{(0)})$ , is often not that large. The use of *conjugate directions* provides a method with a global minimization property.

# Steepest descent and conjugate gradient



(a) Worst case scenario of steepest descent



(b) Conjugate gradient method applied to the same problem as in (a)

# Conjugate directions

Let's revisit equation (3) for a general direction  $\mathbf{d}$  (i.e., not necessarily equal to the negative gradient). Assume  $\mathbf{x} = \mathbf{x}^{(k)}$ , and let  $\mathbf{e}^{(k)} = \mathbf{x}^* - \mathbf{x}^{(k)}$  be the error and  $\mathbf{r}^{(k)} = \mathbf{b} - A\mathbf{x}^{(k)} = A\mathbf{e}^{(k)}$  be the residual. Then we can write  $\langle \mathbf{r}^{(k)}, \mathbf{d} \rangle = \langle \mathbf{e}^{(k)}, \mathbf{d} \rangle_A$ , and so for a general search direction  $\mathbf{d}$  with an exact line search, the iterate takes the form  $\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \frac{\langle \mathbf{e}^{(k)}, \mathbf{d} \rangle_A}{\langle \mathbf{d}, \mathbf{d} \rangle_A} \mathbf{d}$ . By subtracting  $\mathbf{x}^*$ , the iterates in terms of the error  $\mathbf{e}^{(k+1)}$  are given by:

$$\mathbf{e}^{(k+1)} = \mathbf{e}^{(k)} - \frac{\langle \mathbf{e}^{(k)}, \mathbf{d} \rangle_A}{\langle \mathbf{d}, \mathbf{d} \rangle_A} \mathbf{d}. \quad (5)$$

Geometrically, this means that  $\mathbf{e}^{(k+1)}$  is the projection of  $\mathbf{e}^{(k)}$  onto the hyperplane that is  $A$ -orthogonal to  $\mathbf{d}$ , i.e., we have

$$\langle \mathbf{e}^{(k+1)}, \mathbf{d} \rangle_A = 0. \quad (6)$$

## Definition 2 (Conjugate directions)

The vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$  are *conjugate* with respect to a symmetric positive definite matrix  $A$  if they are nonzero and  $A$ -orthogonal:  
 $\langle \mathbf{u}, \mathbf{v} \rangle_A := \langle \mathbf{u}, A\mathbf{v} \rangle = 0$ .

# Conjugate gradient - Warm up

## Theorem 3

Let  $\mathbf{d}^{(0)}, \mathbf{d}^{(1)}, \dots, \mathbf{d}^{(n-1)}$  be  $n$  nonzero pairwise conjugate directions, and consider the sequence of iterates

$$\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha_k \mathbf{d}^{(k)}, \quad \alpha_k = \frac{\langle \mathbf{r}^{(k)}, \mathbf{d}^{(k)} \rangle}{\langle \mathbf{d}^{(k)}, A\mathbf{d}^{(k)} \rangle}.$$

Let  $\mathbf{r}^{(k)} = \mathbf{b} - A\mathbf{x}^{(k)}$  be the residual. Then for each  $k = 1, \dots, n$ ,  $\mathbf{r}^{(k)}$  is orthogonal to  $\text{span}\{\mathbf{d}^{(0)}, \dots, \mathbf{d}^{(k-1)}\}$ . In particular  $\mathbf{r}^{(n)} = \mathbf{0}$ .

## Conjugate gradient - Warm up

**Proof.** Since  $\mathbf{r}^{(k)} = A\mathbf{e}^{(k)}$ , it suffices to show that  $\mathbf{e}^{(k)}$  is  $A$ -orthogonal to  $\text{span}\{\mathbf{d}^{(0)}, \dots, \mathbf{d}^{(k-1)}\}$ . The proof is by induction on  $k$ . For  $k = 0$  there is nothing to prove. Assume the statement is true for  $k \geq 0$ , and consider the equation (5) (with  $\mathbf{d} = \mathbf{d}^{(k)}$ ). From the induction hypothesis, and the fact that the  $\mathbf{d}^{(i)}$  are pairwise conjugate directions, we see that  $\mathbf{e}^{(k+1)}$  is  $A$ -orthogonal to  $\mathbf{d}^{(0)}, \dots, \mathbf{d}^{(k-1)}$ . Furthermore, we have already seen in (6) that  $\langle \mathbf{e}^{(k+1)}, \mathbf{d}^{(k)} \rangle_A = 0$ . Thus this shows that  $\mathbf{e}^{(k+1)}$  is  $A$ -orthogonal to  $\mathbf{d}^{(0)}, \dots, \mathbf{d}^{(k)}$  as desired.  $\square$

## Conjugate gradient - Warm up

---

So, if a sequence ( $\mathbf{d}^{(k)}$ ) of conjugate directions is at hand, we have an iterative procedure with good approximation properties.

The ( $A$ -orthogonal) basis of conjugate directions is constructed by  $A$ -orthogonalization of the sequence  $\{\mathbf{r}_0, A\mathbf{r}_0, A^2\mathbf{r}_0, \dots, A^{n-1}\mathbf{r}_0\}$  with  $\mathbf{r}_0 = \mathbf{b} - A\mathbf{x}_0$ . This is done in the way similar to orthogonalization of the monomial sequence  $\{1, x, x^2, \dots, x^{n-1}\}$  using a recurrence relation.



## Remark 4

It is possible to extend the methods for solving  $A\mathbf{x} = \mathbf{b}$  with symmetric positive definite  $A$  to any other matrices by a simple trick. Suppose we want to solve  $B\mathbf{x} = \mathbf{c}$ , where  $B \in \mathbb{R}^{n \times n}$  is nonsingular. We can convert the above system to the symmetric and positive definite setting by defining  $A = B^T B$ ,  $\mathbf{b} = B^T \mathbf{c}$  and then solving  $A\mathbf{x} = \mathbf{b}$  with the conjugate gradient algorithm (or any other method for positive definite  $A$ ).

# The conjugate gradient method

Here it is.

(A) For any initial vector  $\mathbf{x}^{(0)}$ , set  $\mathbf{d}^{(0)} = \mathbf{r}^{(0)} = \mathbf{b} - A\mathbf{x}^{(0)}$ ;

(B) For  $k \geq 0$ , calculate  $\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha_k \mathbf{d}^{(k)}$  and the residual

$$\mathbf{r}^{(k+1)} = \mathbf{r}^{(k)} - \alpha_k A\mathbf{d}^{(k)}, \quad \text{with}$$
$$\alpha_k := \{ \mathbf{r}^{(k+1)} \perp \mathbf{d}^{(k)} \} = \frac{(\mathbf{r}^{(k)}, \mathbf{d}^{(k)})}{(A\mathbf{d}^{(k)}, \mathbf{d}^{(k)})}, \quad k \geq 0. \quad (7)$$

(C) For the same  $k$ , the next conjugate direction is the vector

$$\mathbf{d}^{(k+1)} = \mathbf{r}^{(k+1)} + \beta_k \mathbf{d}^{(k)}, \quad \text{with}$$
$$\beta_k := \{ \mathbf{d}^{(k+1)} \perp A\mathbf{d}^{(k)} \} = -\frac{(\mathbf{r}^{(k+1)}, A\mathbf{d}^{(k)})}{(\mathbf{d}^{(k)}, A\mathbf{d}^{(k)})}, \quad k \geq 0. \quad (8)$$

## Theorem 5 (Properties of CGM)

For every  $m \geq 0$ , the conjugate gradient method has the following properties.

- (1) The linear space spanned by the residuals  $\{\mathbf{r}^{(i)}\}$  is the same as the linear space spanned by the conjugate directions  $\{\mathbf{d}^{(i)}\}$  and it coincides with the space spanned by  $\{A^i \mathbf{r}^{(0)}\}$ :

$$\text{span}\{\mathbf{r}^{(i)}\}_{i=0}^m = \text{span}\{\mathbf{d}^{(i)}\}_{i=0}^m = \text{span}\{A^i \mathbf{r}^{(0)}\}_{i=0}^m.$$

- (2) The residuals satisfy the orthogonality conditions:  
 $(\mathbf{r}^{(m)}, \mathbf{r}^{(i)}) = (\mathbf{r}^{(m)}, \mathbf{d}^{(i)}) = 0$  for  $i < m$ .
- (3) The directions are conjugate (A-orthogonal):  $(\mathbf{d}^{(m)}, \mathbf{d}^{(i)})_A = (\mathbf{d}^{(m)}, A\mathbf{d}^{(i)}) = 0$  for  $i < m$ .

# The CGM – Theoretical aspects

**Proof.** We use induction on  $m \geq 0$ , the assertions being trivial for  $m = 0$ , since  $\mathbf{d}^{(0)} = \mathbf{r}^{(0)}$  and (2)-(3) are void. Therefore, assuming that the assertions are true for some  $m = k$ , we ask if they remain true when  $m = k + 1$ .

(1) Formula (8)

$$\mathbf{d}^{(k+1)} = \mathbf{r}^{(k+1)} + \beta_k \mathbf{d}^{(k)}$$

readily implies that equivalence of the spaces spanned by  $(\mathbf{r}^{(i)})_0^k$  and  $(\mathbf{d}^{(i)})_0^k$ , is preserved when  $k$  is increased to  $k + 1$ . Similarly, from  $\mathbf{r}^{(k+1)} = \mathbf{r}^{(k)} - \alpha_k A \mathbf{d}^{(k)}$  in (7), and from the inductive assumption  $\mathbf{r}^{(k)}, \mathbf{d}^{(k)} \in \text{span}\{A^i \mathbf{r}^{(0)}\}_{i=0}^k$ , it follows that  $\mathbf{r}^{(k+1)} \in \text{span}\{A^i \mathbf{r}^{(0)}\}_{i=0}^{k+1}$ . To see that  $A^{k+1} \mathbf{r}^{(0)} \in \text{span}\{\mathbf{r}^{(i)}\}_{i=0}^{k+1}$ , since  $\alpha_k \neq 0$ , the claim follows by (5) if  $\mathbf{d}^{(k)}$  has a non-zero component from  $A^k \mathbf{r}^{(0)}$ , and if not the claim follows from the induction hypothesis.

**Proof. Cont.** (2) Turning to assertion (2), we need  $\mathbf{r}^{(k+1)} \perp \mathbf{r}^{(i)}$  for  $i \leq k$ , which by (1) is equivalent to

$$\mathbf{r}^{(k+1)} \perp \mathbf{d}^{(i)} \quad \text{for } i \leq k.$$

We have  $\mathbf{r}^{(k+1)} \perp \mathbf{d}^{(k)}$  by the definition of  $\alpha_k$  in (7), so we need

$$\mathbf{r}^{(k+1)} \stackrel{(7)}{=} \mathbf{r}^{(k)} - \alpha_k A\mathbf{d}^{(k)} \perp \mathbf{d}^{(i)} \quad \text{for } i < k,$$

and this follows from the inductive assumptions  $\mathbf{r}^{(k)} \perp \mathbf{d}^{(i)}$  and  $A\mathbf{d}^{(k)} \perp \mathbf{d}^{(i)}$ .

**Proof. Cont.** (3) It remains to justify (3), namely that  $\mathbf{d}^{(k+1)}$  defined in (8) satisfies

$$\mathbf{d}^{(k+1)} \perp \mathbf{Ad}^{(i)} \quad \text{for } i \leq k.$$

The value of  $\beta_k$  in (8) is defined to give  $\mathbf{d}^{(k+1)} \perp \mathbf{Ad}^{(k)}$ , so we need

$$\mathbf{d}^{(k+1)} \stackrel{(8)}{=} \mathbf{r}^{(k+1)} + \beta_k \mathbf{d}^{(k)} \perp \mathbf{Ad}^{(i)} \quad \text{for } i < k.$$

By the inductive hypothesis  $\mathbf{d}^{(k)} \perp \mathbf{Ad}^{(i)}$ , hence it remains to establish that  $\mathbf{r}^{(k+1)} \perp \mathbf{Ad}^{(i)}$  for  $i < k$ . Now, the formula (7) yields  $\mathbf{Ad}^{(i)} = (\mathbf{r}^{(i)} - \mathbf{r}^{(i+1)})/\alpha_i$ , therefore we require the conditions  $\mathbf{r}^{(k+1)} \perp (\mathbf{r}^{(i)} - \mathbf{r}^{(i+1)})$  for  $i < k$ , and they are a consequence of the assertion (2) for  $m = k + 1$  obtained previously.  $\square$

## Corollary 6 (A termination property)

*If the conjugate gradient method is applied in exact arithmetic, then, for any  $\mathbf{x}^{(0)} \in \mathbb{R}^n$ , termination occurs after at most  $n$  iterations.*

*More precisely, termination occurs after at most  $s$  iterations, where  $s = \dim \text{span}\{A^i \mathbf{r}_0\}_{i=0}^{n-1}$  (which can be smaller than  $n$ ).*

# Termination property

---

**Proof.** Assertion (2) of Theorem 5 states that residuals  $(\mathbf{r}^{(k)})_{k \geq 0}$  form a sequence of mutually orthogonal vectors in  $\mathbb{R}^n$ , therefore at most  $n$  of them can be nonzero. Since they also belong to the space  $\text{span}\{A^i \mathbf{r}_0\}_{i=0}^{n-1}$ , their number is bounded by the dimension of that space. □



# The Krylov subspaces

## Definition 7 (The Krylov subspaces)

Let  $A$  be an  $n \times n$  matrix,  $\mathbf{v} \in \mathbb{R}^n$  nonzero, and  $m \in \mathbb{N}$ . The linear space  $K_m(A, \mathbf{v}) := \text{span}\{A^i \mathbf{v}\}_{i=0}^{m-1}$  is called the *m-th Krylov subspace* of  $\mathbb{R}^n$ .

## Theorem 8 (Number of iterations in CGM)

Let  $A > 0$ , and let  $s$  be the number of its distinct eigenvalues. Then, for any  $\mathbf{v}$ ,

$$\dim K_m(A, \mathbf{v}) \leq s \quad \forall m. \quad (9)$$

Hence, for any  $A > 0$ , the number of iterations of the CGM for solving  $A\mathbf{x} = \mathbf{b}$  is bounded by the number of distinct eigenvalues of  $A$ .

# The Krylov subspaces

**Proof.** Inequality (9) is true not just for positive definite  $A > 0$ , but for any  $A$  with  $n$  linearly independent eigenvectors ( $\mathbf{u}_i$ ). Indeed, in that case one can expand  $\mathbf{v} = \sum_{i=1}^n a_i \mathbf{u}_i$ , and then group together eigenvectors with the same eigenvalues: for each  $\lambda_\nu$  we set  $\mathbf{w}_\nu = \sum_{k=1}^{m_\nu} a_{i_k} \mathbf{u}_{i_k}$  if  $A \mathbf{u}_{i_k} = \lambda_\nu \mathbf{u}_{i_k}$ . Then

$$\mathbf{v} = \sum_{\nu=1}^s c_\nu \mathbf{w}_\nu, \quad c_\nu \in \{0, 1\},$$

hence  $A^i \mathbf{v} = \sum_{\nu=1}^s c_\nu \lambda_\nu^i \mathbf{w}_\nu$ , thus for any  $m$  we get  $K_m(A, \mathbf{v}) \subseteq \text{span}\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_s\}$ , and that proves (9). By Corollary 6, the number of iteration in CGM is bounded by  $\dim K_m(A, \mathbf{r}^{(0)})$ , hence the final conclusion. □

## Remark 9

Theorem 8 shows that, unlike other iterative schemes, the conjugate gradient method is both iterative and direct: each iteration produces a reasonable approximation to the exact solution, and the exact solution itself will be recovered after  $n$  iterations at most.

# Simplifying the CGM-algorithm

We now simplify and reformulate the CGM-algorithm.

Firstly, we rewrite expressions for the parameters  $\alpha_k$  and  $\beta_k$  in (7)-(8) as follows:

$$\alpha_k = \frac{(\mathbf{r}^{(k)}, \mathbf{d}^{(k)})}{(\mathbf{d}^{(k)}, \mathbf{A}\mathbf{d}^{(k)})} \stackrel{(c)}{=} \frac{\|\mathbf{r}^{(k)}\|^2}{(\mathbf{d}^{(k)}, \mathbf{A}\mathbf{d}^{(k)})} > 0,$$
$$\beta_k = -\frac{(\mathbf{r}^{(k+1)}, \mathbf{A}\mathbf{d}^{(k)})}{(\mathbf{d}^{(k)}, \mathbf{A}\mathbf{d}^{(k)})} \stackrel{(a)}{=} -\frac{(\mathbf{r}^{(k+1)}, \mathbf{r}^{(k+1)} - \mathbf{r}^{(k)})}{(\mathbf{d}^{(k)}, \mathbf{r}^{(k+1)} - \mathbf{r}^{(k)})} \stackrel{(b)}{=} \frac{\|\mathbf{r}^{(k+1)}\|^2}{(\mathbf{d}^{(k)}, \mathbf{r}^{(k)})} \stackrel{(c)}{=} \frac{\|\mathbf{r}^{(k+1)}\|^2}{\|\mathbf{r}^{(k)}\|^2} > 0.$$

Here, for  $\beta$ , we used in (a) the fact that  $\mathbf{A}\mathbf{d}^{(k)}$  is a multiple of  $\mathbf{r}^{(k+1)} - \mathbf{r}^{(k)}$  by (7), and in (b) orthogonality of  $\mathbf{r}^{(k+1)}$  to both  $\mathbf{r}^{(k)}$ ,  $\mathbf{d}^{(k)}$  proved in Theorem 5(2). Then, for both  $\beta$  and  $\alpha$ , we used in (c) the property  $(\mathbf{d}^{(k)}, \mathbf{r}^{(k)}) = \|\mathbf{r}^{(k)}\|^2$  which follows from (8) with index  $k + 1$ , taking in account orthogonality  $\mathbf{r}^{(k+1)} \perp \mathbf{d}^{(k)}$ .

Secondly, we let  $\mathbf{x}^{(0)}$  be the zero vector.

# Standard form of the conjugate gradient method

Here it is.

- (1) Set  $k = 0$ ,  $\mathbf{x}^{(0)} = \mathbf{0}$ ,  $\mathbf{r}^{(0)} = \mathbf{b}$ , and  $\mathbf{d}^{(0)} = \mathbf{r}^{(0)}$ ;
- (2) Calculate the matrix-vector product  $\mathbf{v}^{(k)} = A\mathbf{d}^{(k)}$  and  $\alpha_k = \|\mathbf{r}^{(k)}\|^2 / (\mathbf{d}^{(k)}, \mathbf{v}^{(k)}) > 0$ ;
- (3) Apply the formulae  $\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha_k \mathbf{d}^{(k)}$  and  $\mathbf{r}^{(k+1)} = \mathbf{r}^{(k)} - \alpha_k \mathbf{v}^{(k)}$ ;
- (4) Stop if  $\|\mathbf{r}^{(k+1)}\|$  is acceptably small;
- (5) Set  $\mathbf{d}^{(k+1)} = \mathbf{r}^{(k+1)} + \beta_k \mathbf{d}^{(k)}$ , where  $\beta_k = \|\mathbf{r}^{(k+1)}\|^2 / \|\mathbf{r}^{(k)}\|^2 > 0$ ;
- (6) Increase  $k \rightarrow k + 1$  and go back to (2).

# Standard form of the conjugate gradient method

---

The total work is dominated by the number of iterations, multiplied by the time it takes to compute  $\mathbf{v}^{(k)} = A\mathbf{d}^{(k)}$ . Thus the conjugate gradient algorithm is highly suitable when most of the elements of  $A$  are zero, i.e. when  $A$  is *sparse*.