

# Numerical Analysis - Part II

Anders C. Hansen

Lecture 22

---

# *Eigenvalues and eigenvectors*

# Motivation – The Schrödinger equation

---

One of the world's most famous eigenvalue problems: The time independent Schrödinger equation

$$\left[ \frac{-\hbar^2}{2m} \nabla^2 + V(r) \right] \Psi(r) = E\Psi(r).$$

# Introduction to matrix eigenvalue calculations

---

Let  $A$  be a real  $n \times n$  matrix. The eigenvalue equation is  $A\mathbf{w} = \lambda\mathbf{w}$ , where  $\lambda$  is a scalar, which may be complex if  $A$  is not symmetric. There exists a nonzero vector  $\mathbf{w} \in \mathbb{R}^n$  satisfying this equation if and only if  $\det(A - \lambda I) = 0$ . The function  $p(\lambda) = \det(A - \lambda I)$ ,  $\lambda \in \mathbb{C}$ , is a polynomial of degree  $n$ , but calculating the eigenvalues by finding the roots of  $p$  is a disaster area because of loss of accuracy due to rounding errors.

# Introduction to matrix eigenvalue calculations

---

If the polynomial has some multiple roots and if  $A$  is not symmetric, then the number of linearly independent eigenvectors may be fewer than  $n$ , but there are always  $n$  mutually orthogonal real eigenvectors in the symmetric case.

We assume in all cases, however, that the eigenvalue equations  $A\mathbf{w}_i = \lambda_i\mathbf{w}_i$ ,  $i = 1..n$ , are satisfied by eigenvectors  $\mathbf{w}_i$  that are linearly independent, which can be achieved by making an arbitrarily small change to  $A$  if necessary.

# The power method

---

The iterative algorithms that will be studied for the calculation of eigenvalues and eigenvectors are all closely related to the power method, which has the following basic form for generating a single eigenvalue and eigenvector of  $A$ .

We pick a nonzero vector  $\mathbf{x}^{(0)}$  in  $\mathbb{R}^n$ . Then, for  $k = 0, 1, 2, \dots$ , we let  $\mathbf{x}^{(k+1)}$  be a nonzero multiple of  $A\mathbf{x}^{(k)}$ , typically to satisfy  $\|\mathbf{x}^{(k+1)}\| = 1$  so that

$$\mathbf{x}^{(k+1)} = A\mathbf{x}^{(k)} / \|A\mathbf{x}^{(k)}\|, \quad k = 0, 1, 2, \dots$$

This method is oriented on finding an eigenvector corresponding to the largest eigenvalue as the the following theorem shows.

# The power method – Theoretical results

## Theorem 1

Let  $A\mathbf{w}_i = \lambda_i\mathbf{w}_i$ , where the eigenvalues of  $A$  satisfy  $|\lambda_1| \leq \dots \leq |\lambda_{n-1}| < |\lambda_n|$  and the eigenvectors are of the unit length  $\|\mathbf{w}_i\| = 1$ . Assume  $\mathbf{x}^{(0)} = \sum_{i=1}^n c_i \mathbf{w}_i$  with  $c_n \neq 0$ . Then  $\mathbf{x}^{(k)} \rightarrow \pm \mathbf{w}_n$  as  $k \rightarrow \infty$ .

**Proof.** Given  $\mathbf{x}^{(0)}$  as in the assumption,  $\mathbf{x}^{(k)}$  is a multiple of

$$A^k \mathbf{x}^{(0)} = \sum_{i=1}^n c_i \lambda_i^k \mathbf{w}_i = c_n \lambda_n^k \left( \mathbf{w}_n + \sum_{i=1}^{n-1} \frac{c_i}{c_n} \left( \frac{\lambda_i}{\lambda_n} \right)^k \mathbf{w}_i \right).$$

Since  $\|\mathbf{x}^{(k)}\| = \|\mathbf{w}_n\| = 1$ , we conclude that  $\mathbf{x}^{(k)} = \pm \mathbf{w}_n + \mathcal{O}(\rho^k)$ , where the sign is that of  $c_n \lambda_n^k$  and the ratio  $\rho = \frac{|\lambda_{n-1}|}{|\lambda_n|} < 1$  characterizes the rate of convergence. □

# The power method in algorithmic form

Here are the details of an implementation of the procedure.

0. Pick  $\mathbf{x}^{(0)} \in \mathbb{R}^n$  satisfying  $\|\mathbf{x}^{(0)}\| = 1$ . Let  $\varepsilon$  be a small positive tolerance. Set  $k = 0$ .
1. Calculate  $\tilde{\mathbf{x}}^{(k+1)} = A\mathbf{x}^{(k)}$  and set  $\lambda = \frac{\mathbf{x}^{(k)T} A \mathbf{x}^{(k)}}{\mathbf{x}^{(k)T} \mathbf{x}^{(k)}}$ .  
(This  $\lambda$  is called the *Raleigh quotient* and it minimizes  $f(\mu) = \|\tilde{\mathbf{x}}^{(k+1)} - \mu\mathbf{x}^{(k)}\|$  over  $\mu$ .)
2. If  $f(\lambda) \leq \varepsilon$ , accept  $\lambda$  as an eigenvalue and  $\mathbf{x}^{(k)}$  as the corresponding eigenvector.
3. Otherwise, let  $\mathbf{x}^{(k+1)} = \tilde{\mathbf{x}}^{(k+1)} / \|\tilde{\mathbf{x}}^{(k+1)}\|$ , increase  $k$  by one and go back to **1**.



# The power method – Halting criterion

The termination occurs because, by the previous theorem, we have

$$\begin{aligned}\|\tilde{\mathbf{x}}^{(k+1)} - \lambda \mathbf{x}^{(k)}\| &= \min_{\mu} \|\tilde{\mathbf{x}}^{(k+1)} - \mu \mathbf{x}^{(k)}\| \leq \|\tilde{\mathbf{x}}^{(k+1)} - \lambda_n \mathbf{x}^{(k)}\| \\ &= \|A \mathbf{x}^{(k)} - \lambda_n \mathbf{x}^{(k)}\| = \|A \mathbf{w}_n - \lambda_n \mathbf{w}_n\| + \mathcal{O}(\rho^k) = \mathcal{O}(\rho^k) \rightarrow 0.\end{aligned}$$

## Deficiencies of the power method

The power method may perform adequately if  $c_n \neq 0$  and  $|\lambda_{n-1}| < |\lambda_n|$ , where we are using the notation of Theorem 1, but often it is unacceptably slow. The difficulty of  $c_n = 0$  is that, theoretically, in this case the method should find an eigenvector  $\mathbf{w}_m$  with the largest  $m$  such that  $c_m \neq 0$ , but practically computer rounding errors can introduce a small nonzero component of  $\mathbf{w}_n$  into the sequence  $\mathbf{x}^{(k)}$ , and then  $\mathbf{w}_n$  may be found eventually, but one has to wait for the small component to grow.

Moreover,  $|\lambda_{n-1}| = |\lambda_n|$  is not uncommon when  $A$  is real and nonsymmetric, because the spectral radius of  $A$  may be due to a complex conjugate pair of eigenvalues. Next, we will study the inverse iterations (with *shifts*), because they can be highly useful, particularly in the more efficient methods for eigenvalue calculations that will be considered later.

# Inverse iteration

---

This method is highly useful in practice. It is similar to the power method, except that, instead of  $\mathbf{x}^{(k+1)}$  being a multiple of  $A\mathbf{x}^{(k)}$ , we make the choice

$$(A - sI)\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)}, \quad k = 0, 1, \dots, \quad (1)$$

where  $s$  is a scalar that may depend on  $k$  and  $\|\mathbf{x}^{(k)}\| = 1$ . Therefore the calculation of  $\mathbf{x}^{(k+1)}$  from  $\mathbf{x}^{(k)}$  requires the solution of an  $n \times n$  system of linear equations whose matrix is  $(A - sI)$ . Further, if  $s$  is a constant and if  $A - sI$  is nonsingular, we deduce from (1) that  $\mathbf{x}^{(k)}$  is a multiple of  $(A - sI)^{-k}\mathbf{x}^{(0)}$ .

## Inverse iteration

We again let  $\mathbf{x}^{(0)} = \sum_{i=1}^n c_i \mathbf{w}_i$ , as in the proof of Theorem 1, assuming that  $\mathbf{w}_i$ ,  $i = 1..n$ , are linearly independent eigenvectors of  $A$  that satisfy  $A\mathbf{w}_i = \lambda_i \mathbf{w}_i$ . Therefore we note that the eigenvalue equation implies  $(A - sI)\mathbf{w}_i = (\lambda_i - s)\mathbf{w}_i$ , which in turn implies  $(A - sI)^{-1}\mathbf{w}_i = (\lambda_i - s)^{-1}\mathbf{w}_i$ . It follows that  $\mathbf{x}^{(k)}$  is a multiple of

$$(A - sI)^{-k} \mathbf{x}^{(0)} = \sum_{i=1}^n c_i (A - sI)^{-k} \mathbf{w}_i = \sum_{i=1}^n c_i (\lambda_i - s)^{-k} \mathbf{w}_i.$$

Thus, if the  $m$ -th number in the set  $\{|\lambda_i - s|\}$  is the smallest and if  $c_m$  is nonzero, then  $\mathbf{x}^{(k)}$  tends to be a multiple of  $\mathbf{w}_m$  as  $k \rightarrow \infty$ .

We see that the speed of convergence can be excellent if  $s$  is very close to  $\lambda_m$ . Further, it can be made even faster by adjusting  $s$  during the calculation. Typical details are given in the following implementation.

## Typical implementation of inverse iteration

0. Set  $s$  to an estimate of an eigenvalue of  $A$ . Prescribe  $\mathbf{x}^{(0)} \neq 0$ , let  $0 < \varepsilon \ll 1$  and set  $k = 0$ .
1. Calculate (with pivoting if necessary) the LU factorization of  $A - sI$ .
2. Stop if  $U$  is singular because then  $s$  is an eigenvalue of  $A$ , while its eigenvector is any vector in the null space of  $U$ : it can be found easily,  $U$  being upper triangular.
3. Calculate  $\mathbf{x}^{(k+1)}$  by solving  $(A - sI)\mathbf{x}^{(k+1)} = LU\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)}$  using the LU factorization from **1**.
4. Set  $\eta$  to the number that minimizes  $f(\mu) = \|\mathbf{x}^{(k)} - \mu\mathbf{x}^{(k+1)}\|$ .
5. Stop if  $f(\eta) \leq \varepsilon\|\mathbf{x}^{(k+1)}\|$ . Since  $f(\eta) = \|A\mathbf{x}^{(k+1)} - (s + \eta)\mathbf{x}^{(k+1)}\|$ , we let  $s + \eta$  be the calculated eigenvalue of  $A$  and  $\mathbf{x}^{(k+1)}/\|\mathbf{x}^{(k+1)}\|$  be its eigenvector.
6. Otherwise, replace  $\mathbf{x}^{(k+1)}$  by  $\mathbf{x}^{(k+1)}/\|\mathbf{x}^{(k+1)}\|$ , increase  $k$  by one, and either return to **3** without changing  $s$  or to **1** after replacing  $s$  by  $s + \eta$ .

# Upper Hessenberg matrix

---

A matrix of the following form

$$H_n = \begin{bmatrix} h_{1,1} & h_{1,2} & h_{1,3} & \cdots & h_{1,n} \\ h_{2,1} & h_{2,2} & h_{2,3} & \cdots & h_{2,n} \\ 0 & h_{3,2} & h_{3,3} & \cdots & h_{3,n} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & h_{n,n-1} & h_{n,n} \end{bmatrix}.$$

is called an upper Hessenberg matrix.

## Further on inverse iteration

---

The inverse iteration algorithm is very efficient if  $A$  is an *upper Hessenberg matrix*: every element of  $A$  under the first subdiagonal is zero (i.e.  $a_{ij}=0$  if  $j < i-1$ ). In this case the LU factorization in **1** requires just  $\mathcal{O}(n^2)$  or  $\mathcal{O}(n)$  operations when  $A$  is nonsymmetric or symmetric, respectively.

Thus the replacement of  $s$  by  $s + \eta$  in **6** need not be expensive, so fast convergence can often be achieved easily. There are standard ways of giving  $A$  this convenient form which will be considered later.

# Spectrum preserved under similarity transforms

## Theorem 2

Let  $A$  and  $S$  be  $n \times n$  matrices,  $S$  being nonsingular. Then  $\mathbf{w}$  is an eigenvector of  $A$  with eigenvalue  $\lambda$  if and only if  $\hat{\mathbf{w}} = S\mathbf{w}$  is an eigenvector of  $\hat{A} = SAS^{-1}$  with the same eigenvalue.

**Proof.**

$$A\mathbf{w} = \lambda\mathbf{w} \Leftrightarrow AS^{-1}(S\mathbf{w}) = \lambda\mathbf{w} \Leftrightarrow (SAS^{-1})(S\mathbf{w}) = \lambda(S\mathbf{w}).$$

□



# Deflation

Suppose that we have found one solution of the eigenvector equation  $A\mathbf{w} = \lambda\mathbf{w}$ , where  $A$  is again  $n \times n$ . Then *deflation* is the task of constructing an  $(n-1) \times (n-1)$  matrix,  $B$  say, whose eigenvalues are the other eigenvalues of  $A$ . Specifically, we apply a similarity transformation  $S$  to  $A$  such that the first column of  $\hat{A} = SAS^{-1}$  is  $\lambda$  times the first coordinate vector  $\mathbf{e}_1$ , because it follows from the characteristic equation for eigenvalues and from Theorem 2 that we can let  $B$  be the bottom right  $(n-1) \times (n-1)$  submatrix of  $\hat{A} = SAS^{-1}$ . In particular,

$$SAS^{-1} = \hat{A} = \begin{bmatrix} \lambda & \beta \\ 0 & B \end{bmatrix}.$$

We write the condition on  $S$  as  $(SAS^{-1})\mathbf{e}_1 = \lambda\mathbf{e}_1$ . Then the last equation in the proof of Theorem 2 shows that it is sufficient if  $S$  has the property  $S\mathbf{w} = c\mathbf{e}_1$ , where  $c$  is any nonzero scalar.

## Algorithm for deflation for symmetric $A$

---

Suppose that  $A$  is symmetric and  $\mathbf{w} \in \mathbb{R}^n$ ,  $\lambda \in \mathbb{R}$  are given so that  $A\mathbf{w} = \lambda\mathbf{w}$ . We seek a nonsingular matrix  $S$  such that

$$S\mathbf{w} = c\mathbf{e}_1$$

and such that  $SAS^{-1}$  is also symmetric. The last condition holds if  $S$  is orthogonal, since then  $S^{-1} = S^T$ . It is suitable to pick a *Householder reflection*, which means that  $S$  has the form

$$H_{\mathbf{u}} = I - 2\mathbf{u}\mathbf{u}^T / \|\mathbf{u}\|^2, \quad \text{where } \mathbf{u} \in \mathbb{R}^n.$$

## Algorithm for deflation for symmetric $A$

Specifically, we recall from the Numerical Analysis IB course that Householder reflections are orthogonal and that, because  $H_u \mathbf{u} = -\mathbf{u}$  and  $H_u \mathbf{v} = \mathbf{v}$  if  $\mathbf{u}^T \mathbf{v} = 0$ , they reflect any vector in  $\mathbb{R}^n$  with respect to the  $(n-1)$ -dimensional hyperplane orthogonal to  $\mathbf{u}$ . So, for any two vectors  $\mathbf{x}$  and  $\mathbf{y}$  of equal lengths,

$$H_u \mathbf{x} = \mathbf{y}, \quad \text{where} \quad \mathbf{u} = \mathbf{x} - \mathbf{y}.$$

Hence,

$$\left( I - 2 \frac{\mathbf{u}\mathbf{u}^T}{\|\mathbf{u}\|^2} \right) \mathbf{w} = \pm \|\mathbf{w}\| \mathbf{e}_1, \quad \text{where} \quad \mathbf{u} = \mathbf{w} \mp \|\mathbf{w}\| \mathbf{e}_1.$$

Since the bottom  $n-1$  components of  $\mathbf{u}$  and  $\mathbf{w}$  coincide, the calculation of  $\mathbf{u}$  requires only  $\mathcal{O}(n)$  computer operations. Further, the calculation of  $SAS^{-1}$  can be done in only  $\mathcal{O}(n^2)$  operations, taking advantage of the form  $S = I - 2\mathbf{u}\mathbf{u}^T/\|\mathbf{u}\|^2$ , even if all the elements of  $A$  are nonzero.

## Algorithm for deflation for symmetric $A$

---

After deflation, we may find an eigenvector,  $\hat{\mathbf{w}}$  say, of  $SAS^{-1}$ . Then the new eigenvector of  $A$ , according to Theorem 2, is  $S^{-1}\hat{\mathbf{w}} = S\hat{\mathbf{w}}$ , because Householder matrices, like all symmetric orthogonal matrices, are *involutions*:  $S^2 = I$ .



# Givens rotations

---

- 1) We can choose  $\Omega^{[i,j]}$  so that any prescribed element  $\tilde{a}_{jk}$  in the  $j$ -th row of  $\tilde{A} = \Omega^{[i,j]}A$  is zero.
- 2) The rows of  $\tilde{A} = \Omega^{[i,j]}A$  are the same as the rows of  $A$ , except that the  $i$ -th and  $j$ -th rows of the product are linear combinations of the  $i$ -th and  $j$ -th rows of  $A$ .
- 3) The columns of  $\hat{A} = \tilde{A}\Omega^{[i,j]T}$  are the same as the columns of  $\tilde{A}$ , except that the  $i$ -th and  $j$ -th columns of  $\hat{A}$  are linear combinations of the  $i$ -th and  $j$ -th columns of  $\tilde{A}$ .
- 4)  $\Omega^{[i,j]}$  is an orthogonal matrix, thus  $\hat{A} = \Omega^{[i,j]}A\Omega^{[i,j]T}$  inherits the eigenvalues of  $A$ .
- 5) If  $A$  is symmetric, then so is  $\hat{A}$ .

# Transformation to upper Hessenberg – Givens

**Transformation to an upper Hessenberg form:** We replace  $A$  by  $\hat{A} = SAS^{-1}$ , where  $S$  is a product of Givens rotations  $\Omega^{[i,j]}$  chosen to annihilate subsubdiagonal elements  $a_{j,i-1}$  in the  $(i-1)$ -st column:

$$\begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{bmatrix} \xrightarrow{\Omega^{[2,3]} \times} \begin{bmatrix} * & * & * & * \\ \bullet & \bullet & \bullet & \bullet \\ 0 & \bullet & \bullet & \bullet \\ * & * & * & * \end{bmatrix} \xrightarrow{\times \Omega^{[2,3]T}} \begin{bmatrix} * & \bullet & \bullet & * \\ * & \bullet & \bullet & * \\ 0 & \bullet & \bullet & * \\ * & \bullet & \bullet & * \end{bmatrix} \xrightarrow{\Omega^{[2,4]} \times} \begin{bmatrix} * & * & * & * \\ \bullet & \bullet & \bullet & \bullet \\ 0 & * & * & * \\ 0 & \bullet & \bullet & \bullet \end{bmatrix} \xrightarrow{\times \Omega^{[2,4]T}} \begin{bmatrix} * & \bullet & \bullet & \bullet \\ * & \bullet & \bullet & \bullet \\ 0 & \bullet & \bullet & \bullet \\ 0 & \bullet & * & \bullet \end{bmatrix} \xrightarrow{\Omega^{[3,4]} \times} \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ 0 & \bullet & \bullet & \bullet \\ 0 & 0 & \bullet & \bullet \end{bmatrix} \xrightarrow{\times \Omega^{[3,4]T}} \begin{bmatrix} * & \bullet & \bullet & \bullet \\ * & \bullet & \bullet & \bullet \\ 0 & * & \bullet & \bullet \\ 0 & 0 & \bullet & \bullet \end{bmatrix}$$

The  $\bullet$ -elements have changed through a single transformation while the  $*$ -elements remained the same.

It is seen that every element that we have set to zero remains zero, and the final outcome is indeed an upper Hessenberg matrix. If  $A$  is symmetric then so will be the outcome of the calculation, hence it will be tridiagonal. In general, the cost of this procedure is  $\mathcal{O}(n^3)$ .



# Transformation to upper Hessenberg – Householder

Alternatively, we can transform  $A$  to upper Hessenberg using *Householder reflections*, rather than Givens rotations. In that case we deal with a column at a time, taking  $\mathbf{u}$  such that, with  $H_u = I - 2\mathbf{u}\mathbf{u}^T / \|\mathbf{u}\|^2$ , the  $i$ -th column of  $\tilde{B} = H_u B$  is consistent with the upper Hessenberg form. Such a  $\mathbf{u}$  has its first  $i$  coordinates vanishing, therefore  $\hat{B} = \tilde{B}H_u^T$  has the first  $i$  columns unchanged, and all new and old zeros (which are in the first  $i$  columns) stay untouched.

$$\begin{array}{c} \left[ \begin{array}{cccc} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{array} \right] \xrightarrow{H_1 \times} \left[ \begin{array}{cccc} * & * & * & * \\ * & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{array} \right] \xrightarrow{\times H_1^T} \left[ \begin{array}{cccc} * & \bullet & \bullet & \bullet \\ * & \bullet & \bullet & \bullet \\ 0 & \bullet & \bullet & \bullet \\ 0 & \bullet & \bullet & \bullet \\ 0 & \bullet & \bullet & \bullet \end{array} \right] \xrightarrow{H_2 \times} \left[ \begin{array}{cccc} * & * & * & * \\ * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \end{array} \right] \xrightarrow{\times H_2^T} \left[ \begin{array}{cccc} * & * & \bullet & \bullet \\ * & * & \bullet & \bullet \\ 0 & * & \bullet & \bullet \\ 0 & 0 & \bullet & \bullet \\ 0 & 0 & \bullet & \bullet \end{array} \right] \xrightarrow{H_3 \times} \left[ \begin{array}{cccc} * & * & * & * \\ * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{array} \right] \xrightarrow{\times H_3^T} \left[ \begin{array}{cccc} * & * & \bullet & \bullet \\ * & * & \bullet & \bullet \\ 0 & * & \bullet & \bullet \\ 0 & 0 & \bullet & \bullet \\ 0 & 0 & \bullet & \bullet \end{array} \right]
 \end{array}$$

# The QR algorithm

The “plain vanilla” version of the QR algorithm is as follows. Set  $A_0 = A$ . For  $k = 0, 1, \dots$  calculate the QR factorization  $A_k = Q_k R_k$  (here  $Q_k$  is  $n \times n$  orthogonal and  $R_k$  is  $n \times n$  upper triangular) and set  $A_{k+1} = R_k Q_k$ . The eigenvalues of  $A_{k+1}$  are the same as the eigenvalues of  $A_k$ , since we have

$$A_{k+1} = R_k Q_k = Q_k^{-1} (Q_k R_k) Q_k = Q_k^{-1} A_k Q_k, \quad (2)$$

a similarity transformation. Moreover,  $Q_k^{-1} = Q_k^T$ , therefore if  $A_k$  is symmetric, then so is  $A_{k+1}$ .

If for some  $k \geq 0$  the matrix  $A_{k+1}$  can be regarded as “deflated”, i.e. it has the block form

$$A_{k+1} = \begin{bmatrix} B & C \\ D & E \end{bmatrix},$$

where  $B, E$  are square and  $D \approx 0$ , then we calculate the eigenvalues of  $B$  and  $E$  separately (again, with QR, except that there is nothing to calculate for  $1 \times 1$  and  $2 \times 2$  blocks). As it turns out, such a “deflation” occurs surprisingly often.

# The QR iteration for upper Hessenberg matrices

If  $A_k$  is upper Hessenberg, then its QR factorization by means of the Givens rotations produces the matrix

$$R_k = Q_k^T A_k = \Omega^{[n-1,n]} \dots \Omega^{[2,3]} \Omega^{[1,2]} A_k,$$

which is upper triangular. The QR iteration sets

$A_{k+1} = R_k Q_k = R_k \Omega^{[1,2]T} \Omega^{[2,3]T} \dots \Omega^{[n-1,n]T}$ , and it follows that  $A_{k+1}$  is also upper Hessenberg, because

$$\begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \xrightarrow{\times \Omega^{[1,2]T}} \begin{bmatrix} \bullet & \bullet & * & * \\ \bullet & \bullet & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \xrightarrow{\times \Omega^{[2,3]T}} \begin{bmatrix} * & \bullet & \bullet & * \\ * & \bullet & \bullet & * \\ 0 & \bullet & \bullet & * \\ 0 & 0 & 0 & * \end{bmatrix} \xrightarrow{\times \Omega^{[3,4]T}} \begin{bmatrix} * & * & \bullet & \bullet \\ * & * & \bullet & \bullet \\ 0 & * & \bullet & \bullet \\ 0 & 0 & \bullet & \bullet \end{bmatrix}$$

Thus a strong advantage of bringing  $A$  to the upper Hessenberg form initially is that then, in every iteration in QR algorithm,  $Q_k$  is a product of just  $n-1$  Givens rotations. Hence each iteration of the QR algorithm requires just  $\mathcal{O}(n^2)$  operations.