

## Example Sheet 1

1. Consider the two-dimensional flow  $u = 1/(1+t)$ ,  $v = 1$  in  $t > -1$ . Find and sketch
- the streamline at  $t = 0$  that passes through the point  $(1, 1)$ ,
  - the path of a fluid particle released from  $(1, 1)$  at  $t = 0$ ,
  - the position at  $t = 0$  of a streak of dye released from  $(1, 1)$  during the time interval  $-1 < t \leq 0$ .

2. A steady two-dimensional flow (pure straining) is given by  $u = \alpha x$ ,  $v = -\alpha y$  with  $\alpha$  constant.

- Find the equation for a general streamline of the flow, and sketch some of them.
- At  $t = 0$  the fluid on the curve  $x^2 + y^2 = a^2$  is marked (by an electro-chemical technique). Find the equation for this material fluid curve for  $t > 0$ .
- Does the area within the curve change in time, and why?

3. Repeat question 2 for the two-dimensional flow (simple shear) given by  $u = \gamma y$ ,  $v = 0$  with  $\gamma$  constant. Which of the two flows stretches the curve faster at long times?

4. A two-dimensional flow is represented by a streamfunction  $\psi(x, y)$  with  $u = \partial\psi/\partial y$  and  $v = -\partial\psi/\partial x$ . Show that

- the streamlines are given by  $\psi = \text{const}$ ,
- $|\mathbf{u}| = |\nabla\psi|$ , so that the flow is faster where the streamlines are closer,
- the volume flux per unit  $z$ -distance crossing any curve from  $\mathbf{x}_0$  to  $\mathbf{x}_1$  is given by  $\psi(\mathbf{x}_1) - \psi(\mathbf{x}_0)$ ,
- $\psi = \text{const}$  on any *fixed* (i.e. stationary) boundary.  
[Hint for (iii):  $\mathbf{n} ds = (dy, -dx)$ .]

5. Verify that the two-dimensional flow given in Cartesian coordinates by

$$u = \frac{y-b}{(x-a)^2 + (y-b)^2}, \quad v = \frac{a-x}{(x-a)^2 + (y-b)^2}$$

satisfies  $\nabla \cdot \mathbf{u} = 0$ , and then find the streamfunction  $\psi(x, y)$  such that  $u = \partial\psi/\partial y$  and  $v = -\partial\psi/\partial x$ . Sketch the streamlines.

6. Verify that the two-dimensional flow given in polar coordinates by

$$u_r = U \left( 1 - \frac{a^2}{r^2} \right) \cos \theta, \quad u_\theta = -U \left( 1 + \frac{a^2}{r^2} \right) \sin \theta$$

satisfies  $\nabla \cdot \mathbf{u} = 0$ , and find the streamfunction  $\psi(r, \theta)$ . Sketch the streamlines.

$$\left[ \text{Take: } \nabla \cdot \mathbf{u} = \frac{1}{r} \frac{\partial}{\partial r} (ru_r) + \frac{1}{r} \frac{\partial}{\partial \theta} (u_\theta) \quad \text{and} \quad u_r = \frac{1}{r} \frac{\partial \psi}{\partial \theta}, \quad u_\theta = -\frac{\partial \psi}{\partial r} \right]$$

7. Verify that the axisymmetric flow (uniaxial straining) given in cylindrical polar coordinates by  $u_r = -\frac{1}{2}\alpha r$ ,  $u_z = \alpha z$  satisfies  $\nabla \cdot \mathbf{u} = 0$ , and find the Stokes streamfunction  $\Psi(r, z)$ . Sketch the streamlines.

$$\left[ \text{Take: } \nabla \cdot \mathbf{u} = \frac{1}{r} \frac{\partial}{\partial r} (r u_r) + \frac{\partial u_z}{\partial z} \quad \text{and} \quad u_r = -\frac{1}{r} \frac{\partial \Psi}{\partial z}, \quad u_z = \frac{1}{r} \frac{\partial \Psi}{\partial r} \right]$$

8. An axisymmetric jet of water of speed  $1 \text{ m s}^{-1}$  and cross-section  $6 \times 10^{-4} \text{ m}^2$  strikes a wall at right angles and spreads out over it. By using the momentum integral equation, calculate the force on the wall due to the jet. [Neglect gravity.]

9. Starting from the Euler momentum equation for a fluid of constant density with a potential force  $-\nabla\Phi$  per unit mass, show that for fixed volume  $V$  enclosed by surface  $A$

$$\frac{d}{dt} \int_V \frac{1}{2} \rho |\mathbf{u}|^2 dV + \int_A \rho H \mathbf{u} \cdot \mathbf{n} dA = 0,$$

where  $H$  is the Bernoulli quantity,  $H = \frac{1}{2} |\mathbf{u}|^2 + \frac{p}{\rho} + \Phi$ . (This is sometimes interpreted as saying that  $H$  is the ‘transportable energy’, or the ‘advected energy’, per unit mass.)

10. How high can water rise up one’s arm hanging in the river from a lazy ( $1 \text{ m s}^{-1}$ ) punt? [Use Bernoulli on surface streamline, where  $p = 1$  atmosphere.]

11. A flat-bottomed barge closely fits a canal, so that while it travels very slowly it still generates a fast  $5 \text{ m s}^{-1}$  current under it. Estimate how much lower in the water the barge lies as a result of this current. [*Hint*: Archimedes when stationary. Flow reduces pressure, so have to go deeper to get same pressure on long bottom.]

12. Waste water flows into a large tank at  $10^{-4} \text{ m}^3 \text{ s}^{-1}$  and out of a short exit pipe of cross-section  $4 \times 10^{-5} \text{ m}^2$  into the air. The flow has reached a steady state. Estimate the height of the free surface of the water in the tank, relative to the height of the exit pipe.

13. A water clock is an axisymmetric vessel with a small exit pipe in the bottom. Find the shape of vessel for which the water level falls equal heights in equal intervals of time.