

# Hyperbolic equations

Avoid numerically

▶ Advection + diffusion

OK if  $\Delta x < D/U$ . Then  $D\Delta t < \Delta x^2$  gives  $U\Delta t < \Delta x$

▶ Advection + reaction

OK if  $\Delta x < U\tau$ . Then  $U\Delta t < \Delta x$  gives  $\Delta t < \tau$


▶ Pure Advection

- ▶ Problem 1 conserve past numerical errors
- ▶ Problem 2 shocks = unresolved boundary layers = rarefaction waves and discontinuities ← unfriendly to high-order schemes


Hint: Reformulate with characteristics, i.e. Lagrangian

## 1.1 Simplest - unstable

First-order in time, central second-order in space

$$\frac{u_\ell^{n+1} - u_\ell^n}{\Delta t} = -c \frac{u_{\ell+1}^n - u_{\ell-1}^n}{2\Delta x}$$


$ct = 0.0 (0.2) 1.0$   
 $\Delta x = 0.05$   
 $c\Delta t = 0.0125$



Unstable

# 1. Simple smooth advection

$$u_t + cu_x = 0,$$

and smooth initial condition

$$u(x) = \begin{cases} 4(x-1)^2(2-x)^2 & \text{in } 1 \leq x \leq 2, \\ 0 & \text{otherwise.} \end{cases}$$

Take  $c$  constant,  $> 0$ .

Generalise to  $c(x)$ ,  $c(x, u)$  and vector  $\mathbf{u}(\mathbf{x}, t)$

Finite differences easier for cooperation of spatial and temporal discretisations.

Write

$$u_\ell^n = u(x = \ell\Delta x, t = n\Delta t).$$

## Stability analysis

Set  $u_\ell^n = A^n e^{ik\ell\Delta x}$ , (Fourier wave). To find  $A(k)$

Algorithm  $\rightarrow A = 1 - i\mu \sin \theta$  with  $\mu = \frac{c\Delta t}{\Delta x}$  and  $\theta = k\Delta x$ .

Then  $|A| > 1$  all  $\mu$ ,

i.e. **unstable** all  $\Delta t$ .


Most unstable = short wave zigzag  $\theta = \frac{\pi}{2}$

with  $|A| = \sqrt{1 + \mu^2}$

i.e.  $u \sim (1 + \mu^2)^{t/2\Delta t}$ .

## 1.2 Lax-Friedricks – too stable

Replacing  $u_\ell^n$  in the time derivative by average  $\frac{1}{2}(u_{\ell+1}^n + u_{\ell-1}^n)$ .



$$u_\ell^{n+1} = \frac{1}{2} \left( 1 - \frac{c\Delta t}{\Delta x} \right) u_{\ell+1}^n + \frac{1}{2} \left( 1 + \frac{c\Delta t}{\Delta x} \right) u_{\ell-1}^n.$$

Stability analysis  $u_\ell^n = A^n e^{ik\ell\Delta x}$

$$A = \cos \theta - i\mu \sin \theta \quad \text{with } \mu = \frac{c\Delta t}{\Delta x} \text{ and } \theta = k\Delta x$$


i.e. stable  $|A| < 1$  all  $\theta$  if

$$\mu = \frac{c\Delta t}{\Delta x} < 1 \quad \text{CFL condition (Courant-Friedricks-Lewy)}$$


Information propagates less than  $\Delta x$  in  $\Delta t$

## Lax-Friedricks – too stable

Plots  $ct = 0.0 (0.2) 1.0$ ,  $\Delta x = 0.05$



unstable  $\mu = c\Delta t/\Delta x = 1.1$



stable  $\mu = 0.5$

Stable but very damped

## Longwave error analysis

Taylor series


$$u_{\ell+1}^n = u_\ell^n + \Delta x u_x \Big|_\ell^n + \frac{1}{2} \Delta x^2 u_{xx} \Big|_\ell^n + \dots,$$

$$u_\ell^{n+1} = u_\ell^n + \Delta t u_t \Big|_\ell^n + \frac{1}{2} \Delta t^2 u_{tt} \Big|_\ell^n + \dots$$

Algorithm + Lax trick  $u_{tt} = c^2 u_{xx}$

$$u_t = -cu_x + \frac{1}{2}(1 - \mu^2) \frac{\Delta x^2}{\Delta t} u_{xx}.$$

Numerical diffusion




$ct = 0.0 (0.2) 1.0$   
for  $\Delta x = 0.05$   
continuous  $c\Delta t = 0.025$   
dashed  $c\Delta t = 0.0125$

NB numerical diffusion  $\nearrow$  as  $\Delta t \rightarrow 0$

## 1.3 Upwinding – avoid downstream influence

$$\frac{u_\ell^{n+1} - u_\ell^n}{\Delta t} = -c \frac{u_\ell^n - u_{\ell-1}^n}{\Delta x}$$




Stability

$$|A|^2 = 1 - 4\mu(1 - \mu) \sin^2 \frac{\theta}{2}, \quad \text{i.e. stable if } \mu < 1$$

Longwave error analysis

$$u_t = -cu_x + \frac{1}{2}(1 - \mu)c\Delta x u_{xx}.$$




$ct = 0.0 (0.2) 1.0$   
 $\Delta x = 0.05$ ,  $\Delta t = 0.25$

numerical diffusivity bounded  
as  $\Delta t \rightarrow 0$

## 1.4 Crank-Nicolson – second-order, implicit

Central difference about mid-point  $(\ell, n + \frac{1}{2})$



$$\frac{u_\ell^{n+1} - u_\ell^n}{\Delta t} = -\frac{c\Delta t}{4\Delta x} (u_{\ell+1}^{n+1} - u_{\ell-1}^{n+1} + u_{\ell+1}^n - u_{\ell-1}^n).$$

Stability

$$A = \frac{1 - \frac{1}{2}i\mu \sin \theta}{1 + \frac{1}{2}i\mu \sin \theta}.$$

i.e.  $|A| = 1$  all  $\mu$ : **stable** with no damping (?accurate large  $\mu$ ?)

## 1.5 Lax-Wendroff – second-order, explicit

Upwinding corrected by subtracting off leading error

$$\frac{1}{2}(1 - \mu)c\Delta x [u_{xx} \approx (u_{\ell+1}^n - 2u_\ell^n + u_{\ell-1}^n)/\Delta x^2]$$

and rearranging

$$u_\ell^{n+1} = u_\ell^n - \frac{c\Delta t}{2\Delta x} (u_{\ell+1}^n - u_{\ell-1}^n) + \frac{c^2\Delta t^2}{2\Delta x^2} (u_{\ell+1}^n - 2u_\ell^n + u_{\ell-1}^n)$$

Stability

$$|A|^2 = 1 - 4\mu^2(1 - \mu^2)\sin^4 \frac{1}{2}\theta,$$

**stable** if  $\mu < 1$  (CFL)

Longwave errors

$$u_t = -cu_x - \frac{1}{6}(1 - \mu^2)c\Delta x^2 u_{xxx}.$$


again **numerical dispersion**

## Crank-Nicolson

Longwave error analysis

$$u_t = -cu_x - \frac{1}{12}(2 - \mu^2)c\Delta x^2 u_{xxx}.$$

$u_{xxx}$  means **numerical dispersion**




$ct = 0.0(0.2) 1.0$   
 $\Delta x = 0.05, c\Delta t = 0.025$

Slower short waves at the trailing edge

## Lax-Wendroff

$$u_\ell^{n+1} = u_\ell^n - \frac{c\Delta t}{2\Delta x} (u_{\ell+1}^n - u_{\ell-1}^n) + \frac{c^2\Delta t^2}{2\Delta x^2} (u_{\ell+1}^n - 2u_\ell^n + u_{\ell-1}^n)$$




$ct = 0.0(0.2) 1.0$   
 $\Delta x = 0.05, c\Delta t = 0.025$

Slower short waves at the trailing edge

## 1.6 Angled derivative – second-order, explicit, 3-level

Central difference about mid-point  $(\ell - \frac{1}{2}, n)$




$$(u_t)_{\ell-\frac{1}{2}}^n = \frac{1}{2} \left( \frac{u_{\ell-1}^n - u_{\ell-1}^{n-1}}{\Delta t} + \frac{u_{\ell}^{n+1} - u_{\ell}^n}{\Delta t} \right) = -c(u_x)_{\ell-\frac{1}{2}}^n = c \frac{u_{\ell}^n - u_{\ell-1}^n}{\Delta x}.$$

Re-arranging

$$u_{\ell}^{n+1} = \left( 1 - \frac{2c\Delta t}{\Delta x} \right) (u_{\ell}^n - u_{\ell-1}^n) + u_{\ell-1}^{n-1}.$$

## Angled derivative

$$\text{Start } u_{\ell}^1 = u_{\ell}^0 - \frac{c\Delta t}{2\Delta x} (u_{\ell+1}^0 - u_{\ell-1}^0)$$



$$ct = 0.0 \ (0.2) \ 1.0$$

$$\mu = 0.3$$

## Angled derivative

Stability

$$\left( Ae^{i\theta/2} \right)^2 - 2i(1 - 2\mu) \sin \frac{1}{2}\theta \left( Ae^{i\theta/2} \right) - 1 = 0,$$

stable  $\mu < 1$ , but spurious (stable) **second mode**

Longwave errors

$$u_t = -cu_x + \frac{1}{12}(1 - \mu)(1 - 2\mu)c\Delta x^2 u_{xxx}.$$

**numerical dispersion**, vanishes at  $\mu = \frac{1}{2}$  (when exact!)

## Conclusions for smooth problems

CFL stability:  $\mu = \frac{c\Delta t}{\Delta x} < 1$  (typically)

Odd-order schemes  $\rightarrow$  **numerical diffusion**

i.e. spreading and decay

Even-order schemes  $\rightarrow$  **numerical dispersion**

i.e. spurious (typically trailing) oscillations