

Comments and corrections to acla2@damtp.cam.ac.uk.

1. Let $u, v \in \mathcal{D}'(\mathbf{R}^n)$, one of which has compact support. Show that the convolution $u * v$, as defined in your notes, is uniquely defined and gives rise to an element of $\mathcal{D}'(\mathbf{R}^n)$.
2. Show that if $u, v, w \in \mathcal{D}'(\mathbf{R}^n)$ and at least two of them have compact support, then the convolution is associative, i.e. $(u * v) * w = u * (v * w)$.
3. Let $\varphi \in \mathcal{D}(\mathbf{R})$ and choose $\epsilon > 0$ sufficiently small so that $\text{supp}(\varphi) \subset I_\epsilon$, where $I_\epsilon = (-1/\epsilon, 1/\epsilon)$. Given that φ has a uniformly convergent Fourier series on I_ϵ in the form

$$\varphi(x) = \sum_{n \in \mathbf{Z}} c_n e^{i\epsilon\pi n x}, \quad c_n = \frac{\epsilon}{2} \int \varphi(x) e^{-i\epsilon\pi n x} dx,$$

prove the Fourier inversion theorem on $\mathcal{D}(\mathbf{R})$ by taking a suitable limit.

4. For $\varphi \in \mathcal{S}(\mathbf{R}^n)$ prove that $\sum_m \varphi(m) = \sum_n \hat{\varphi}(2\pi n)$. This is the famous Poisson summation formula.
5. If $u \in H^s(\mathbf{R}^n)$ show that $D^\alpha u \in H^{s-|\alpha|}(\mathbf{R}^n)$. If $s > t$ show that $H^s(\mathbf{R}^n) \subset H^t(\mathbf{R}^n)$.
6. Show that $\mathcal{S}(\mathbf{R}^n)$ is dense in $L^2(\mathbf{R}^n) = H^0(\mathbf{R}^n)$ and deduce that $\mathcal{S}(\mathbf{R}^n)$ is dense in $H^s(\mathbf{R}^n)$, i.e. prove that for each $u \in H^s(\mathbf{R}^n)$ there is a sequence $\{\varphi_m\}_{m \geq 1}$ in $\mathcal{S}(\mathbf{R}^n)$ such that

$$\lim_{m \rightarrow \infty} \|u - \varphi_m\|_{H^s} = 0.$$

[Hint: Use Parseval's theorem.]

7. Prove that multiplication by a Schwartz function gives rise to a continuous map from $H^s(\mathbf{R}^n)$ to itself, i.e. $\|\varphi u\|_{H^s} \lesssim \|u\|_{H^s}$ for $\varphi \in \mathcal{S}(\mathbf{R}^n)$. You may assume Peetre's inequality: for $\lambda, \mu \in \mathbf{R}^n$ and $s \in \mathbf{R}$

$$\left(\frac{1 + |\lambda|^2}{1 + |\mu|^2} \right)^s \leq 2^{|s|} (1 + |\lambda - \mu|^2)^{|s|}.$$

8. For $u \in \mathcal{S}'(\mathbf{R}^n)$ show that $(D^\alpha u)^\wedge = \lambda^\alpha \hat{u}$ and $(x^\beta u)^\wedge = (-D)^\beta \hat{u}$ for all multi-indices α, β .
9. For $u \in \mathcal{E}'(\mathbf{R}^n)$ show that $\hat{u}(\lambda) = \langle u(x), e^{-i\lambda \cdot x} \rangle$. Deduce that if $u \in \mathcal{E}'(\mathbf{R}^n)$ there exists some $t \in \mathbf{R}$ such that $u \in H^t(\mathbf{R}^n)$.
10. Suppose that $u_1, u_2 \in \mathcal{S}'(\mathbf{R}^n)$ and at least one of u_1 and u_2 has compact support. Show that their convolution $u_1 * u_2 \in \mathcal{S}'(\mathbf{R}^n)$ and $(u_1 * u_2)^\wedge = \hat{u}_1 \hat{u}_2$.
11. Show that $e^{-\epsilon x} H \rightarrow H$ in $\mathcal{S}'(\mathbf{R})$ as $\epsilon \rightarrow 0$. Hence show

$$\hat{H} = \pi \delta_0 - \text{ip.v.} \left(\frac{1}{x} \right)$$

in $\mathcal{S}'(\mathbf{R})$.

12. The Riemann-Lebesgue lemma states that if $u \in L^1(\mathbf{R})$ then $|\hat{u}(\lambda)| \rightarrow 0$ as $|\lambda| \rightarrow \infty$. Prove this result by considering the substitution $x = x' + \pi/\lambda$ in the integral defining $\hat{u}(\lambda)$.
13. If $u \in H^m(\mathbf{R}^n)$ with $m \in \mathbf{N}$, use Parseval's theorem to show that

$$\sum_{|\alpha| \leq m} \int |D^\alpha u|^2 dx < \infty.$$

Prove the converse.

14. Let $\mathcal{O}(\mathbf{R}^n)$ denote the space of smooth functions that grow no faster than a polynomial. Show that $\mathcal{O}(\mathbf{R}^n) \subset \mathcal{S}'(\mathbf{R}^n)$. Fix $\varphi \in \mathcal{S}(\mathbf{R}^n)$ with $\varphi(0) = 1$. For $u \in \mathcal{O}(\mathbf{R}^n)$ we define

$$\hat{u}_\epsilon(\lambda) = \int e^{-i\lambda \cdot x} \varphi(\epsilon x) u(x) dx.$$

Show that $\hat{u} = \lim_{\epsilon \rightarrow 0} \hat{u}_\epsilon$ in $\mathcal{S}'(\mathbf{R}^n)$.

15. Compute the Fourier transform in $\mathcal{S}'(\mathbf{R})$ of the function

$$u(x) = \frac{x}{1+x^2}.$$

For which $s \in \mathbf{R}$ is $u \in H^s(\mathbf{R})$?

16. Prove that $D^\alpha \delta_0 \in H^s(\mathbf{R}^n)$ if and only if $s < -|\alpha| - \frac{1}{2}n$.

17. Let $\Gamma = \{x \in \mathbf{R}^n : x \cdot n = 0\}$ be a hyperplane with surface element $d\sigma$ and normal n . Let $\chi \in \mathcal{S}(\mathbf{R}^n)$ be a fixed Schwartz function and set $d\mu = \chi d\sigma$. This defines a distribution $\mu_\Gamma \in \mathcal{S}'(\mathbf{R}^n)$ by

$$\langle \mu_\Gamma, \varphi \rangle = \int_\Gamma \varphi d\mu$$

for each $\varphi \in \mathcal{S}(\mathbf{R}^n)$. Prove that

$$\hat{\mu}_\Gamma(\lambda) = \int_\Gamma e^{-i\lambda \cdot x} d\mu(x).$$

Classify the large $|\lambda|$ behaviour of $\hat{\mu}_\Gamma$. For which $s \in \mathbf{R}$ is $\mu_\Gamma \in H^s(\mathbf{R}^n)$?

18. Suppose that $u \in \mathcal{S}'(\mathbf{R}^n)$ and $\Delta^2 u + u \in H^s(\mathbf{R}^n)$. Prove that $u \in H^{s+4}(\mathbf{R}^n)$.

19. Compute the Fourier transforms of the functions

$$(a) \operatorname{sgn}(x), \quad (b) \arctan(x), \quad (c) x \log|x| - x, \quad (d) \exp(i\omega x^2)$$

in $\mathcal{S}'(\mathbf{R})$, where $\omega \in \mathbf{R}$.

20. Show that the Fourier transform of $(x_1 + ix_2)^{-1}$ is proportional to itself and find the constant of proportionality.

21. Suppose $f \in C(\mathbf{R})$ and $f(x) - 1/x = O(1/x^2)$ as $|x| \rightarrow \infty$. Prove that

$$\lim_{\epsilon \rightarrow 0} \left[\hat{f}(-\epsilon) - \hat{f}(+\epsilon) \right] = 2\pi i.$$

22. If $\varphi \in \mathcal{D}(\mathbf{R}^n)$ and $\operatorname{supp}(\varphi) \subset B_\delta$ show that $\hat{\varphi}(z)$ is an entire function and there exist constants C_m such that

$$|\hat{\varphi}(z)| \leq C_m (1 + |z|)^{-m} e^{\delta |\operatorname{Im} z|}$$

for $m = 0, 1, 2, \dots$ and $z \in \mathbf{C}^n$. Prove the converse.