EXAMPLES II

1. Vacuum energy density In special and general relativity, the energy density ρ and pressure P of a perfect fluid combine to form the *energy-momentum* tensor. In the rest frame of the fluid, this is given by

$$T^{\mu\nu} = \begin{bmatrix} \rho & 0 & 0 & 0 \\ 0 & P & 0 & 0 \\ 0 & 0 & P & 0 \\ 0 & 0 & 0 & P \end{bmatrix} .$$

After a Lorentz boost by velocity v in the $x-$direction, the energy-momentum tensor transforms to

$$\widetilde{T}^{\mu\nu} = \Lambda^{\mu}{}_{\lambda}\Lambda^{\nu}{}_{\sigma}T^{\lambda\sigma} \quad \text{with} \quad \Lambda^{\mu}{}_{\lambda} = \left[\begin{array}{cc|cc} \gamma & -\frac{\gamma\, v}{c} & 0 & 0 \\ -\frac{\gamma\, v}{c} & \gamma & 0 & 0 \\ \hline 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right] \quad \text{and} \quad \gamma = \frac{1}{\sqrt{1-\frac{v^2}{c^2}}} .$$

What equation of state $P = \omega\,\rho$ gives rise to a fluid that looks the same to all inertial observers?

2. Deceleration parameter is defined by

$$q(t) \equiv -\frac{a\ddot{a}}{\dot{a}^2} .$$

Compute q for a flat universe dominated by a single fluid with equation of state $P = \omega\rho$.

For a universe with arbitrary curvature, filled with matter, radiation and a cosmological constant, show that today

$$q_0 = \frac{1}{2}\Omega_{\rm m} + \Omega_{\rm rad} - \Omega_\Lambda .$$

What is the deceleration parameter for our universe?

Show that the Taylor expansion of the scale factor about the present day (with $a(t_0) = 1$) can be written as

$$\frac{1}{1+z} = a(t) = 1 + H_0(t-t_0) - \frac{1}{2}q_0H_0^2(t-t_0)^2 + \dots .$$

Hence show that the light from a nearby galaxy at redshift $z \ll 1$ was emitted at time

$$t_0 - t = H_0^{-1}z - \frac{1}{2}H_0^{-1}(2+q_0)z^2 + \dots .$$

3. Matter–dark energy transition. A flat universe, containing matter and a cosmological constant, expands according to the Friedmann equation

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3c^2}\left(\rho_\Lambda + \frac{\rho_0}{a^3}\right) .$$

Solve this to find

$$a(t) = \beta(\sinh\,\alpha t)^{2/3} \tag{1}$$

with α and β constants that you should specify. [Hint: you may find the substitution $b = a^{3/2}$ helpful.] Verify the expected asymptotic behaviour for early and late times. smallskip

Our universe today is flat, with $\Omega_{\rm m} \approx 0.3$ and $\Omega_\Lambda \approx 0.7$ and $H_0^{-1} \approx 14\,$Gyr. Using these parameters, estimate:

- The age of the universe.
- The age at which the expansion of the universe first started to accelerate.
- The age at which the energy density in matter and the cosmological constant were equal.

A cosmologically ignorant civilisation measures $H_0^{-1} \approx 14\,$Gyr but refuses to countenance the possibility of a cosmological constant. How old would they believe their universe to be if they assumed it was flat with $\Omega_{\rm m} = 1$?

Copyright © 2023 University of Cambridge. Not to be quoted or reproduced without permission.

4. Beyond expansion and acceleration. The *jerk* (or *surge*) parameter is defined as

$$J(t) = \frac{\dddot{a}\, a^2}{\dot{a}^3}\,.$$

The past 14 billion years of cosmic expansion can be roughly characterised as: the jerk is unity. To see this, consider the Friedmann equation with matter, a cosmological constant and curvature. Show that the curvature can be written as

$$k\, c^2 = a^2 H^2 (J - 1)\,.$$

Confirm that the solution (1) indeed has unit jerk. (Note k is defined to take values: $k = +1/R_0^2, 0, -1/R_0^2$.) [Fun fact: the 4th, 5th and 6th derivatives of the scale factor are called *snap*, *crackle* and *pop* respectively.]

5. Radiation–Λ model. A flat universe, containing radiation and a cosmological constant, expands according to the Friedmann equation

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3c^2}\left(\rho_\Lambda + \frac{\rho_0}{a^4}\right)\,.$$

Solve for $a(t)$. Verify the expected asymptotic behaviour for early and late times.

6. Virial theorem and dark matter. Consider a collection of particles, each of which interacts through the potential

$$V(\mathbf{x}_i - \mathbf{x}_j) = \alpha\, |\mathbf{x}_i - \mathbf{x}_j|^n\,.$$

Prove the virial theorem,

$$2\overline{T} = n\,\overline{V}\,,$$

where $\overline{T}$ is the time-averaged kinetic energy and $\overline{V}$ the time-averaged total potential energy. Show that conventional orbits in our solar system satisfy this relation.

7. Exact inflationary solution. The equations of motion for inflation can be written as

$$H^2 = \frac{1}{3\,M_{\text{pl}}^2}\left[\frac{1}{2}\dot{\phi}^2 + V(\phi)\right] \tag{2}$$

and

$$\ddot{\phi} + 3H\dot{\phi} + \frac{\partial V}{\partial \phi} = 0\,, \tag{3}$$

where $M_{\text{pl}}^2 = c^2/(8\pi G)$ is related to the *reduced Planck mass*. Consider the potential

$$V(\phi) = V_0\, e^{-\frac{\lambda\phi}{M_{\text{pl}}}}$$

Show that the inflationary equations have the exact solution

$$a(t) = \left(\frac{t}{t_0}\right)^{\frac{2}{\lambda^2}} \quad \text{and} \quad \phi(t) = \phi_0 + \alpha \log t\,,$$

for some choice of ϕ_0 and α.

For what values of λ does inflation occur? Define the pressure-energy density ratio $\omega_\phi = P_\phi/\rho_\phi$. For what range of ω_ϕ does inflation occur?

8. Hamilton–Jacobi approach to inflation. Assume that an inflationary solution $\phi(t)$ is monotonic and view the Hubble parameter $H(t)$ as a function $H(\phi(t))$. Show that the equations of inflation (2) and (3) can be written in "Hamilton-Jacobi" form

$$\dot{\phi} = -2M_{\text{pl}}^2 \frac{\partial H}{\partial \phi}$$

and

$$V(\phi) = 3M_{\text{pl}}^2 H^2 - 2M_{\text{pl}}^4 \left(\frac{\partial H}{\partial \phi}\right)^2\,.$$

[Hint: you might start by differentiating the Friedmann equation (2) with respect to time.]

Copyright © 2023 University of Cambridge. Not to be quoted or reproduced without permission.

The Hamilton-Jacobi formalism can be used to construct exact solutions in which one specifies a choice of $H(\phi)$ and then works backwards to read off the corresponding potential $V(\phi)$, the scalar evolution $\phi(t)$ and, hence the scale factor $a(t)$. Find these three quantities for

$$H(\phi) = \exp\left(-\sqrt{\frac{\alpha}{2}}\frac{\phi}{M_{\rm pl}}\right) .$$

9. Slow-roll approximation. Under the slow roll conditions, the inflationary equations become

$$H^2 \approx \frac{1}{3M_{\rm pl}^2}V(\phi) \quad \text{and} \quad 3H\dot{\phi} \approx -\frac{\partial V}{\partial \phi} .$$

Solve these equations to find the scale factor $a(t)$ for the choice of potentials

- $V(\phi) = \frac{1}{2}m^2\phi^2$;
- $V(\phi) = \frac{1}{4}\lambda\phi^4$.

Determine the value of ϕ when inflation ends and hence the initial ϕ_0 required to achieve 60 e-folds.

**Please send any corrections to epss@damtp.cam.ac.uk*

Copyright © 2023 University of Cambridge. Not to be quoted or reproduced without permission.