

### Examples Sheet 1

1. (Warm-up) Writing the amplitude for a quantum mechanical particle, mass  $m$ , to freely propagate in 1-dimension as

$$\langle x | e^{-iH(t-t_0)} | x_0 \rangle \equiv K(x, t; x_0, t_0) = \sqrt{\frac{m}{2\pi i(t-t_0)}} \exp\left(\frac{im(x-x_0)^2}{2(t-t_0)}\right)$$

with  $t > t_0$ , show that

$$\int dx' K(x, t; x', t') K(x', t'; x_0, t_0) = K(x, t; x_0, t_0)$$

with  $t > t' > t_0$ . Verify that  $\lim_{t \rightarrow 0} K(x, t; 0, 0) = \delta(x)$  and that

$$i \frac{\partial}{\partial t} K(x, t; 0, 0) = -\frac{1}{2m} \frac{\partial^2}{\partial x^2} K(x, t; 0, 0).$$

Use these facts to express the solution of the Schrödinger equation for a free particle,  $\Psi(x, t)$  in terms of an initial wavefunction  $\Psi(x, 0)$  and  $K(x, t; 0, 0)$ . Check this result for  $\Psi(x, 0) = e^{ikx}$ , with  $k$  constant.

2. Consider the quantum mechanics of a particle moving in 1-dimension with Hilbert space  $\mathcal{H}$ . Obtain path integral expressions, in imaginary time  $T$ , for the following,

(a)  $\text{Tr}_{\mathcal{H}}(\mathbf{P}e^{-TH})$ , where  $\mathbf{P}$  is the parity operator  $\mathbf{P} : x \mapsto -x$ , and the trace of an operator  $O$ ,  $\text{Tr}_{\mathcal{H}}(O)$ , is the sum or integral over the expectation values of  $O$  in a complete basis of states.

(b)  $\langle \psi_f | e^{-TH} | \psi_i \rangle$ , where  $\psi_{i,f}(x) = \langle x | \psi_{i,f} \rangle$  are arbitrary states in the Hilbert space.

For a particle in a 1-dimensional harmonic potential  $V(x) = \frac{1}{2}m\omega^2x^2$ , the amplitude for particle propagation in real time  $t$  is

$$K(x, t; x_0, 0) = \sqrt{\frac{m\omega}{2\pi i \sin \omega t}} \exp\left(im\omega \frac{(x^2 + x_0^2) \cos \omega t - 2xx_0}{2 \sin \omega t}\right)$$

(e.g. see Osborn's notes §1.2.1). Using this amplitude for  $t = -iT$ , evaluate your expressions for (a) and (b) explicitly in the case that  $|\psi_{i,f}\rangle$  are the ground state of the harmonic oscillator. Check that they agree with what you expect from quantum mechanics, working directly in the energy basis.

3. Consider the partition function

$$Z(\lambda) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dx \exp\left(-\frac{1}{2}x^2 - \frac{\lambda}{4!}x^4\right)$$

for a zero-dimensional QFT, given  $\lambda > 0$ .

(a) By expanding the integral in  $\lambda$ , obtain the  $n$ -th order perturbative expression

$$Z_n(\lambda) = \sum_{\ell=0}^n \left(-\frac{\lambda}{4!}\right)^\ell \frac{(4\ell)!}{4^\ell(2\ell)! \ell!} \quad (1)$$

and show that, for  $\ell \leq 3$ , the coefficients  $a_\ell$  of  $\lambda^\ell$  in this expression are the sums of symmetry factors of the relevant loop Feynman graphs. [At 2-loop order there is only 1 graph, at 3-loops there are 2, and at 4-loops there are 4.]

(b) (Optional but instructive.) Using any computer package, plot  $Z_n(\lambda = 0.1)$  against  $n$  to see that there is a region in  $n$  where  $Z_n$  appears to converge before blowing up as  $n$  is increased.

(c) (\* Slightly beyond the scope of the course.) Show that the minimum value of  $a_\ell \lambda^\ell$  occurs when  $\ell \approx \frac{3}{2\lambda}$ . Hence show that the Borel transform

$$\mathcal{B}Z(\lambda) = \sum_{\ell=0}^{\infty} \frac{1}{\ell!} a_\ell \lambda^\ell$$

converges provided  $|\lambda| < \frac{3}{2}$  and that in this case

$$Z(\lambda) = \int_0^\infty dz e^{-z} \mathcal{B}Z(z\lambda)$$

so that  $Z(\lambda)$  may be recovered from its Borel transform.

(d) By expanding  $e^{-x^2/2}$  in the integral in (1) obtain the strong coupling expansion

$$Z(\lambda) = \frac{1}{2\sqrt{\pi}} \sum_{L=0}^{\infty} \frac{(-1)^L}{L!} \Gamma\left(\frac{L}{2} + \frac{1}{4}\right) \left(\frac{6}{\lambda}\right)^{\frac{L}{2} + \frac{1}{4}}$$

for  $Z(\lambda)$  as a series in  $1/\sqrt{\lambda}$ . For  $\lambda = \frac{1}{10}$  how many terms does one need in order to obtain the value at which the weak coupling expansion appeared to converge?

4. Let  $e^{-W(J)/\hbar} = \int d^n \phi e^{-(S(\phi) + J_c \phi_c)/\hbar}$ , and let  $\Gamma(\Phi)$  be the Legendre transform of  $W(J)$ . Show directly that

$$\hbar^2 \frac{\partial^3 W}{\partial J_a \partial J_b \partial J_c} \Big|_{J=0} = \langle \phi_a \phi_b \phi_c \rangle^{\text{conn}}$$

and that

$$-\frac{1}{\hbar} \frac{\partial^3 \Gamma}{\partial \Phi_a \partial \Phi_b \partial \Phi_c} \Big|_{J=0} = \langle \phi_a \phi_b \phi_c \rangle_{\text{1PI}}^{\text{conn}}.$$

5. In lectures we showed that

$$\int d^{2k} \theta e^{\frac{1}{2} A_{ab} \theta_a \theta_b} = \text{Pf}(A)$$

where  $A$  is a real, invertible antisymmetric matrix and  $\theta_a$  are  $2k$  Grassmann variables. By writing  $\theta_a = N_{ab}\theta'_b$  for some real matrix  $N$ , show that  $\text{Pf}(N^T A N) = \det(N)\text{Pf}(A)$ . Show that  $N$  may be chosen so as to put  $A$  into the form

$$N^T A N = \begin{pmatrix} 0 & 1 & & & & \\ -1 & 0 & & & & \\ & & 0 & 1 & & \\ & & -1 & 0 & & \\ & & & & \ddots & \\ & & & & & 0 & 1 \\ & & & & & -1 & 0 \end{pmatrix},$$

and demonstrate that consequently  $\text{Pf}(A) = \pm\sqrt{\det A}$ .

6. Consider a theory of 4 Grassmann variables  $\theta_a$ ,  $a = 1 \dots 4$ , governed by the action

$$S(\theta) = \frac{1}{2} A_{ab} \theta_a \theta_b + \frac{1}{4!} \lambda_{abcd} \theta_a \theta_b \theta_c \theta_d.$$

Compute the partition function of this theory (a) by directly expanding  $e^{-S/\hbar}$  in the path integral and (b) by writing down the Feynman rules and drawing all possible vacuum diagrams.

7. Let  $M$  be an  $N \times N$  Hermitian matrix and let

$$V(M) = \frac{1}{2} \text{Tr} M^2 + \frac{g}{4} \text{Tr} M^4$$

where  $g$  is a constant.

- Show that  $V(M)$  is invariant under  $M \mapsto U^\dagger M U$ , where  $U$  is a unitary matrix. Explain why this implies that  $V(M)$  only depends on the eigenvalues  $\{\lambda_i\}$  of  $M$ .
- Viewing  $V(M)$  as the action for a zero-dimensional QFT, obtain an expression for the propagator  $\langle M_{ij} M_{kl} \rangle$  correct to lowest nontrivial order in  $g$ . [Consider only connected diagrams.]
- Now let  $B$ ,  $C$ , and  $H$  be further  $N \times N$  matrices, each of which have zeros all along the leading diagonal. Let the elements of  $B$  and  $C$  be fermionic variables, while the entries of  $H$  are bosonic. Consider the matrix integral

$$Z(g) = \int \frac{[dM dB dC dH]}{(2\pi i)^{N(N-1)}} \exp[-NV(M) + i \text{Tr}(HM) + \text{Tr}(B[M, C])]$$

where the measure  $[dM dB dC dH]$  indicates an integral over each entry of  $M$  and the off-diagonal entries of  $B$ ,  $C$ , and  $H$ . Obtain the effective action for the eigenvalues  $\{\lambda_i\}$  of  $M$ . [Do not attempt to perform the path integral over these eigenvalues.]

Please e-mail me at [M.Wingate@damtp.cam.ac.uk](mailto:M.Wingate@damtp.cam.ac.uk) with any comments, especially any errors.